The Effect Of Carnosine On Lower Extremity Ischemia Reperfusion İnjury İn Rat

Purpose: The purpose of this study is; to examine the effect of carnosine on ischemia

reperfusion injury in lungs and muscle occuring after occlusion – reperfusion of infrarenal

abdominal aorta in rats.

Material and Method: Twentyfour Sprague-Dawley rats were randomly divided into 3

groups (n=8). Sham laparotomy (control) group underwent laparotomy and dissection of the

infrarenal abdominal aorta without occlusion and saline was given intraperitoneally. In aortic

ischemia reperfusion groups (group 2 and 3), after laparotomy and exteriorization of the

infrarenal abdominal aorta, aorta was occluded for 30 min ischemia and then the clamp was

removed for 60 min reperfusion. In group 2; saline and in group 3; 250mg/kg carnosine, in

equal volumes, were given intraperitoneally 10 min before declamping of the aorta. The

blood, lung and muscle specimens were taken for biochemical analysis of malonil dialdehyde,

superoxide dismutase, catalase, myeloperoxidase, nitric oxide, glutathione peroxidase,

glutathione reductase, total glutathione and oxide glutathione. Additionally, in blood samples

total antioksidant capasity was analyzed.

Results: İn lung specimens, levels of malonil dialdehyde, superoxide dismutase,

myeloperoxidase, oxide glutathione and total glutathione in the aortik ischemia reperfusion +

saline group (group 2) were significantly higher than the levels in the sham laparotomy

(control) group (p<0.05). Level of glutathione reductase, malonil dialdehyde and oxide

glutathione in the aortik ischemia reperfusion + carnosine group (group 3) was significantly

lower than the level in the aortik ischemia reperfusion + saline group (p<0.05).

Conclusion: Carnosine could prevent the systemic and local effects of abdominal aortic

occlusion-reperfusion injury in rats. Since this is a first demonstration for the use of carnosine in

abdominal aortic occlusion-reperfusion injury, the results of the present study could light the

way for the future studies.

Key word: Ischemia Reperfusion Injury, Gastrocnemius, Lung, Carnosine

e-mail: drbilgealacam@yahoo.com.tr